

Norton Sound Summer Commercial Red King Crab Fishery
Observer Project Summary Report, 1992

By

Elisabeth Brennan

Regional Informational Report¹ No. 3A92-22

Alaska Department of Fish and Game
Division of Commercial Fisheries, AYK Region
333 Raspberry Road
Anchorage, Alaska 99518

September 1992

¹ The Regional Information Report Series was established in 1987 to provide an information access system for all unpublished divisional reports. These reports frequently serve diverse ad hoc informational purposes or archive basic uninterpreted data. To accommodate timely reporting of recently collected information, reports in this series undergo only limited internal review and may contain preliminary data; this information may be subsequently finalized and published in the formal literature. Consequently, these reports should not be cited without prior approval of the author or the Division of Commercial Fisheries.

TABLE OF CONTENTS

	<u>Page</u>
TABLE OF CONTENTS.....	i
LIST OF TABLES.....	ii
LIST OF FIGURES.....	v
INTRODUCTION.....	1
Objectives and Tasks.....	2
METHODS.....	2
RESULTS.....	3
Harvest Summary.....	3
Legal Male King Crab.....	4
Sublegal Male King Crab.....	4
Female King Crab.....	5
Bycatch.....	5
Tags.....	6
DISCUSSION.....	6
Harvest Summary.....	6
Legal Male King Crab.....	6
Sublegal Male King Crab.....	7
Female King Crab.....	7
Bycatch.....	8
National Marine Fisheries Service Trawl Survey.....	8
COMMENTS ON OBSERVER PROGRAM.....	8
SUMMARY.....	9
LITERATURE CITED.....	10

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Red king crab verbal catch reports for all fishing vessels by statistical area, Norton Sound Section, Eastern Bering Sea, for August 1, 12pm through August 1, 12 pm through August 3, 8am 1992.....	11
2. Red king crab season total catch (from fish ticket reports) by statistical area for Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	12
3. Observer crab catch summary; number of observed pot lifts, average soak time, number of legal males, sublegal males and females and proportion (%) of each captured by statistical area, and the corresponding mean number of crab caught per observed pot lift (CPP) and pot lift standardized to a 24 hour soak period (CPP/24hr), Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	13
4. Carapace length measurement summary of sampled legal male red king crab captured during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	14
5. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 656401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	15
6. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 666401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	16
7. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 636401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	17
8. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 656330 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	18

LIST OF TABLES (continued)

<u>Table</u>	<u>Page</u>
9. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 666330 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	19
10. Percent of newly recruited male king crab by statistical area, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	20
11. Carapace length measurement summary of Prerecruit, Recruit, and Postrecruit male king crab, captured during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	21
12. Observer summary table of red king crab weight samples by statistical area, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	22
13. Carapace length measurement summary of sampled sublegal male red king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-5, 1992.....	23
14. Carapace length measurement summary of sampled sublegal male red king crab captured in statistical area 656401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	24
15. Carapace length measurement summary of sampled sublegal male red king crab captured in statistical area 666401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	25
16. Carapace length measurement summary of sampled sublegal male red king crab captured in statistical area 636401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	26
17. Carapace length measurement summary of sampled sublegal male red king crab captured in statistical area 656330 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992 (no old shell sample)	27

LIST OF TABLES (continued)

<u>Table</u>	<u>Page</u>
18. Carapace length measurement and percent ovigerity summary of sampled female red king crab captured during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	28
19. Carapace length measurement and percent ovigerity summary of sampled female red king crab captured in statistical area 656401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	29
20. Carapace length measurement and percent ovigerity summary of sampled female red king crab captured in statistical area 666401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	30
21. Carapace length measurement and percent ovigerity summary of sampled female red king crab captured in statistical area 636401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	31

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. King crab fishing districts and sections of Statistical Area Q.....	32
2. Statistical areas for the Norton Sound red king crab fishery.....	33
3. Length frequency distribution and frequency of new and old carapace age condition of legal male red king crab, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	34
4. Length frequency distribution of prerecruit, recruit, and postre- cruit male red king crab, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	35
5. Length frequency distribution and frequency of new and old carapace age condition of prerecruit male king crab, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	36
6. Length frequency distribution of female red king crab, juveniles and adults, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.....	37

INTRODUCTION

The Norton Sound Section of the Northern Bering Sea District consists of all waters in statistical area Q that are north of the latitude of Cape Romanzof, east of 168 west longitude, and south of the latitude of Cape Prince of Wales (Figures 1 and 2). A large vessel summer commercial red king crab (Paralithodes camtschatica) fishery has existed in the Norton Sound Section since 1977. No summer commercial fishery occurred in 1991 due to a lack of staff to properly manage the fishery. The budget had been cut the previous winter.

The National Marine Fisheries Service conducted their triennial trawl survey to examine the abundance of Norton Sound red king crab in late August 1991. The results of that survey as compared to the 6 previous trawl surveys show a gradual trend of increasing abundance since the low of 1982, with 3.4 million pounds of legal king crab in the commercial fishing district in 1991. The quota for the Norton Sound Section for the 1992 season was therefore set at 300,000 pounds.

The 1992 summer commercial red king crab fishery opened at 12 noon, August 1 in the Norton Sound Section of the Eastern Bering Sea. The emergency order closure was also announced at that time for 12 noon, August 3.

Twenty seven vessels took part in the summer commercial crab season. Five of these vessels were catcher/processors and each had an independent fishery observer on board. One ADF&G observer was placed on a catcher vessel. This is the third year subcontracted observers were placed on board commercial vessels in the Norton Sound Section. It is the tenth season an ADF&G observer was placed on board a crab vessel in Norton Sound. In 1989, the Board of Fisheries adopted a regulation (5AAC 39.645) mandating that all catcher/processors and floating processors which process shellfish have independent observers on board. These observers are certified by the ADF&G and work for independent contractors. The onboard observer program provides the only effective means of collecting essential biological and management data from vessels that process shellfish. This data is necessary in determining the magnitude and location of the commercial harvest and the status of the stock. The observers also provide the only effective means to enforce size and sex restriction regulations that will protect the resource. Catcher vessels are not required to have observers onboard, but may choose to allow an ADF&G observer onboard to collect data essential to the management of the fishery.

A new regulation (5AAC 34.925) was put into effect June 19, 1992 regarding pot limit. A king crab pot limit for statistical area Q states that no more than 100 pots may be operated from a vessel.

Objectives and Tasks

The specific objectives and tasks of the observer program are to:

1. Report daily the number of pots pulled and number of legal crab harvested for each statistical area fished by the vessel on which the observer is placed and by any vessels delivering to the observer's vessel.
2. Obtain and record daily 100 length frequency samples of harvested legal male red king crab (carapace width $\geq 4\text{-}3/4$ inches), 100 length frequency samples of incidentally caught sublegal male red king crab, and at least 20 length frequency samples of incidentally caught female red king crab.
3. Determine the mean live weight of the harvested legal male crab.
4. Determine the carapace age of the sampled crab.
5. Determine the percentage of new recruits in the commercial harvest.
6. Determine the degree of ovigerity of sampled female red king crab.
7. Determine the percentage of illegal commercial harvest by sampling a minimum of 600 crab over the course of each day for legal size and sex.
8. Develop a relative abundance index for legal male, sublegal male and female king crab by recording the catch of as many pot lifts as is required to obtain a sample size of 300 or more crab per day.
9. Document bycatch species by number if possible (optional).

METHODS

The methods used during the 1992 commercial crab season for catch reporting, sampling crab, skipper interviews, and collecting information from tagged crab are presented in the ADF&G Shellfish Observer Field Manual and Addendum #1 to Observers Manual (found on the inside back cover of Observer Manual). This manual is available through the ADF&G commercial fishery office in Nome. The identity of vessels from which observer data was collected have been omitted from this report to maintain confidentiality.

Quality and quantity of the different types of data collected varied from observer to observer. Some data had statistical area information omitted. Length frequency data was partially lost by one observer when his briefcase fell into the water on his way to debriefing at the ADF&G office in Nome. One observer collected no bycatch data.

RESULTS

Harvest Summary

Catch reporting logs were kept by observers on each catcher/processor and by skippers of catcher vessels for each statistical area fished. Verbal reports were relayed 8:00 a.m., August 3 to the ADF&G office in Nome. The verbal reports indicated that six statistical areas were fished (Table 1).

Fish ticket reports document that five statistical areas were fished (Table 2). This data showed that this years largest fishing effort and greatest productivity occurred in statistical area 656401 (Table 2). Forty-seven percent of all pot lifts occurred in area 656401, 31% in area 666401, 7% in area 636401, 7% in area 666330, and 7% in area 656330. Area 656401 accounted for 73% of the total harvest, area 666401 for 14%, area 656330 for 6%, area 666330 for 5%, and area 636401 for 2%. Total pots pulled from individual statistical areas were: 2714 pots in area 656401, 1804 pots in area 666401, 419 pots for area 636401, 421 pots in area 656330, and 388 pots in area 666330. Total harvest for individual areas were: 18,193 crab in area 656401, 3351 crab in area 666401, 1646 crab in area 656330, 1344 crab in area 666330, and 368 crab in area 636401. The total number of pots pulled during the season were 5,746 and the total commercial harvest was 24,902 crab. Total poundage was 74,029 pounds of crab.

Catcher Log Summary Sheets were kept by all six observers. Information was collected from four statistical areas and is presented in Table 3. The overall average soak time per pot was 15.3 hours. Observed catch per pot (CPP) of legal male red king crab by area were: 2.2 crab/pot in area 666401, 2.1 crab/pot in area 656330, 2.0 crab/pot in area 656401, and 0.6 crab/pot in area 636401. Overall observed CPP for legal male king crab was 1.9 crab/pot. Observed CCP standardized for a 24 hour soak period (CPP/24hr) for legal males was 3.9 in area 666401, 3.3 in area 656401, 2.4 in area 656330, and 0.7 in area 636401. Catch per pot from fish ticket data were 6.7 crab/pot in area 656401, 3.9 crab/pot in area 656330, 3.5 crab/pot in area 666330, 1.9 crab/pot in area 666401, and 0.9 crab/pot in area 636401. Overall CCP for the season was 4.3 crab/pot (Table 2).

Legal male red king crab made up 92% of the observed catch in area 656330, 75% in area 656401, 71% in area 666401, and 35% in area 636401 (Table 3). Approximately 70% of the overall observed catch were legal male crab. Sublegal male crab made up 23% of the overall observed catch and female crab 7% of the observed catch.

Five of the six observers provided usable information about illegal harvest on their vessels. A total of 3,474 harvested crab were sampled. Three crab, or 0.08%, were found to be illegal. Two of these illegal crab were sublegal males and one was a female blue king crab. The percent of illegal catch by individual vessels ranged from 0% to 0.7% of the harvest.

Legal Male King Crab

Carapace length, age and crab weight were collected by five observers from the catch of each vessel. One observer handed in partial data at debriefing. A few length frequency forms had no statistical area information recorded on them.

Carapace length measurements were taken from 2,566 legal male red king crab in five statistical areas during the extent of the 1992 summer fishery. Carapace age was classified as new (11 months old) or old (at least 23 months old) (Table 4, Figure 3).

Overall mean carapace length of sampled legal male king crab was 119.7 mm. Mean carapace length of sampled legal males for separate statistical areas were: 117.5 mm in area 656401, 120.7 mm in area 666401, 120.9 mm in area 636401, 127.6 mm in area 656330, and 119.3 in area 666330 (Tables 5-9).

Crab with new shell carapace made up 71% of the total legal male king crab sampled while old shell crab made up 29% of the sample (Table 4). New shell crab made up 66.2% of the crab sampled in area 656401, 76% in area 666401, 65% in area 636401, 77% in area 656330, and 67% in area 666330 (Tables 5-9).

Recruit red king crab are defined as new shell, legal male king crab with a carapace length less than or equal to 115 mm. Recruit crab made up 28% of all legal male crab sampled (Table 4). Post recruit crab made up 72% of the legal male crab sampled. The recruit king crab composition ranged from 6 - 46% for individual statistical areas sampled throughout the fishery (Table 10). The total sampled male catch was composed of 13% prerecruits, 25% recruits, and 62% post recruits (Table 11).

The mean weight of legal red king crab was calculated from a sample of 463 individuals. Average weight ranged from 2.4 to 3.7 lbs./crab. The overall mean weight was 3.0 pounds (Table 12).

One purple pigmented legal red king crab was found during sampling. The crab was photographed.

Sublegal (prerecruit) Male King Crab

Carapace length measurement and shell age was collected from a sample of 598 sublegal male red king crab (Table 13, Figure 5). Data was collected from four statistical areas.

The overall mean length of the sublegal crab sampled was 87.7 mm (Table 13). Mean length for area 656401 was 90.9 mm, area 666401 was 87.7 mm, area 636401 73.1 mm, and area 656330 was 94.7 mm (Tables 14-17).

New shell crab made up 91% of the sample and old shell crab made up 9% of the overall sample (Table 13). The sample from area 656401 was made up of 89% new shell crab, area 666401 was made up of 90% new shell crab, and areas 636401 and 656330 were 100% new shell crab (Tables 14-17).

Overall observed mean catch of sublegal male king crab per pot, standardized to a 24 hour soak period (CPP/24hr) was 1 crab/pot (Table 3).

Female King Crab

Carapace length measurement and percent ovigerity was collected from a total of 167 female red king crab during the commercial fishery (Table 18, Figure 6). Data was collected from three statistical areas (Tables 18-21). Mature female king crab made up 77% of all females sampled. Immature female crab made up 23% of the sample. Seventy-three percent of the mature female crab were considered to have a high degree of ovigerity ($\geq 60\%$).

The mean carapace length of mature female crab was 93.5 mm, and 66.8 mm for immature female crab (Table 18). In area 656401, 34 female crab were sampled. Mean carapace length was 66 mm for immature crab and 88.5 mm for mature crab. In area 666401, 92 female crab were sampled. Mean carapace length was 81 mm for immature crab and 97.7 mm for mature crab. In area 636401, 36 female crab were sampled. Mean carapace length was 66.3 mm for immature crab and 75 mm for mature crab (Tables 19-21).

Overall observed mean catch per pot of female king crab standardized to a 24 hour soak period (CPP/24hr) was 0.3 crab/pot (Table 3). All observers reported that most mature female red king crab sampled had dark purple eggs in their clutch.

Bycatch

Information on bycatch was collected to some degree by five of the six observers. Quantifiable data was collected by two observers. Other observers noted number of species or estimated the number without collecting number of pots pulled, soak time, and statistical areas. Quantifiable information collected shows at least 471 pots sampled for a total bycatch yield of 626 opilio tanner crab Chionoecetes opilio, 436 arctic lyre crab Hyas coarctatus, 166 blue king crab (135 female, 19 male, and 12 no sex recorded) Paralithodes platypus, 113 starry flounder Platichthys stellatus, 29 yellowfin sole Limanda aspera, 6 pacific cod Gadus macrocephalus, 5 korean hair crab Erimacrus isenbeckii, 4 pacific halibut Hippoglossus stenolepis, 3 sculpin, and 1 pollock Theragra chalcogramma. Observers recorded nonquantifiable information as "most pots had kelp crab and opilio", "numerous starfish and lyre crab", "lots" of starry flounder, yellowfin sole and sculpin, and "hundreds" of starfish.

One apparent "hybrid" red-blue king crab was found in a pot pulled from area 666401. It was a new shell crab with carapace length of 106 mm. The rostrum resembled that of a blue king crab, and had 3 pairs of prominent spines on the center of the carapace like that of a red king crab. Coloring was two toned (orange/blue) on legs, and the carapace was orange/red with blue/purple spotting. Several photos were taken to record the color and it was bagged and frozen and brought to the ADF&G office in Nome.

Tags

One crab was found with a tag during the commercial fishery. The crab was captured August 2, 1992 in statistical area 656330, in 10 fathoms of water. Only the tag was recovered, no carapace length or age was collected. The tag ID was NX 01,860.

DISCUSSION

Harvest Summary

Five catcher/processor vessels and 22 catcher vessels took part in the 1992 summer commercial red king crab fishery in the Norton Sound Section.

Fish ticket records show area 656401 accounted for 47% of the pot pulls and 73% of the total harvest (Table 2). The total number of crab harvested was 24,902, total pots pulled was 5,746. The total harvest was 74,029 pounds. The harvest goal was 300,000 pounds.

Based on fish ticket reports, area 656401 had a CPP of 6.7, area 666401 had a CPP of 1.9 (Table 2). Overall catch per pot for the season was 4.3 crab/pot. Catch per pot standardized for a 24 hour soak period (CPP/24hr) was 3.0 overall. Observer crab catch summary shows area 656401 with 2.0 crab/pot and area 666401 with 2.2 crab/pot. Overall CPP from the observer summary was 1.9 crab/pot.

Legal male crab made up 75% of the observed catch in area 656401 and 71% of the observed catch in area 666401 (Table 3). These statistical areas are located relatively close to shore. Legal male crab made up 92% of the observed catch in area 656330 which is further offshore (Table 3). Adult male king crab are larger and tend to migrate offshore farther and faster than smaller sublegal male and female crab so this trend is expected. Legal male king crab made up 70% of the overall observed catch. Legal males tend to range farther offshore than sublegal.

Five of the six onboard observers sampled 3,474 crab from the commercial catch to determine the illegal harvest level on individual fishing vessels. Illegal catch levels ranged from 0% to 0.7% of the harvest. The illegal harvest was found to be 0.08% overall. This is well below the 3.0% limit thought to be citable by Fish and Wildlife Protection. The onboard observer program contributes a great deal to this low illegal harvest level.

Legal Male King Crab

The overall mean carapace length of sampled legal male red king crab was 119.7 mm. Sample size was 2566 crab. This is similar to data from previous years. Mean carapace length was 119.0 mm in 1988 (Gebhard and Lean, 1988), 119.8 mm in 1989 (Gebhard and Lean, 1989), and 121.1 mm in 1990 (Gebhard, 1990).

This seasons legal male new shell/old shell ratio was 71% new shell and 29% old shell (Table 4). Legal male new shell/old shell ratio in 1990 was 83% new shell and 17% old shell. In 1989 this ratio was 71% new shell and 29% old shell.

Total recruitment to the harvested stock was 28% of the population (Table 4). Total post recruits made up 72% of the harvested stock. This compares to the 1990 harvest, for which total recruits made up 21% and total post recruits made up 79% of the harvested stock. In 1989 recruits comprised 23% of the harvested stock, and in 1988 they comprised 25%.

In the observed male catch, prerecruits, recruits and post recruits made up 13%, 25%, and 62% of the harvested population (Table 11, Figure 4). In 1990 these figures were 21% prerecruits, 17% recruits, and 62% post recruits.

Mean weight of the harvested legal crab was 3.0 pounds per crab. Samples ranged from 2.4 to 3.7 pounds (Table 12). In 1990 average weight was 3.9 pounds, in 1989 average weight was 3.1 pounds, and in 1988 average weight was 3.2 pounds.

Sublegal (Prerecruit) Male King Crab

The overall mean length of the sampled sublegal male crab was 87.4 mm (Table 13). This is similar to mean lengths of 87.1 mm in 1990 and 88.4 mm in 1989.

New shell crab made up 90.9% of the total sublegal sample (Table 13). This is compared to 88% new shell crab in 1990, 75% in 1989, and 82% in 1988.

Overall observed mean catch of sublegal male red king crab standardized for a 24 hour soak period (CPP/24hr) was 1.0 crab/pot (Table 3). Samples were obtained from area 656401 (1.0 CCP/24hr), area 666401 (1.2 CPP/24hr), area 636401 (0.9 CPP/24hr), and area 656330 (0.2 CPP/24hr).

Female King Crab

Mature female red king crab made up 77% of females sampled throughout the duration of the fishery. Immature female crab made up 23% of the sample. This compares to 18% immature female crab in 1990, and 10% immature female crab in 1989.

Mean carapace length was 93.5 mm for mature female crab and 66.8 mm for immature female crab (Table 18). The sample size was 129 adult female crab and 38 immature female crab. Mean carapace length for adult female crab was 81.2 mm in 1990, 80.1 mm in 1989, and 82.5 mm in 1988. Immature female carapace length was 65.9 mm in 1990, 68.1 mm in 1989, and 67.6 mm in 1988.

A large portion of the mature female crab, 73%, was considered to have a high degree of ovigerity ($\geq 60\%$). This is a decline from previous years; 75% in 1990, 78% in 1989, and 89% in 1988. All observers noted that most mature female crab had dark purple eggs in their clutch. This indicated uneyed eggs. Eyed brown eggs are usually observed at this time of year.

Overall observed mean catch per pot of female red king crab standardized to a 24 hour soak period (CPP/24hr) was 0.3 crab/pot. This compares to 0.2 CPP/24hr in 1990 and 3.2 CPP/24hr in 1989.

Bycatch

Bycatch information was collected thoroughly by two observers, and to some degree by three others. This years bycatch included 11 identified species: opilio tanner crab, arctic lyre crab, blue king crab, starry flounder, yellow fin sole, pacific cod, korean hair crab, halibut, sculpin, pollock, and starfish. At least 471 pots from three different statistical areas were sampled for bycatch. Information was not collected in the systematic way it had been in previous seasons, therefore no estimated bycatch could be calculated.

One apparent "hybrid" red-blue king crab was found in a pot pulled from area 666401. The number of spines and coloring were different from what has been seen in the past.

National Marine Fisheries Service Trawl Survey

The National Marine Fisheries Service (NMFS) conducted their triennial survey of red king crab in Norton Sound August 22 to 30, 1991. Results show length frequency modes at about 70 mm and 110 mm for male crab, and at about 60 mm for female crab. The majority of crabs <80 mm were found to be recently molted. All females were new, hardshell crab. All female crab \geq 80 mm were ovigerous. The NMFS estimated abundance of legal male red king crab in Norton Sound was 1.38 million crab, or 4.0 million pounds, of which 85% or 3.4 million pounds was in the legal fishing district. This compares to 1.78 million pounds in the fishing district in 1988 and 2.2 million pounds in 1985.

Data collected during the 1992 summer commercial fishery show immature king crab were not present as was indicated in the NMFS trawl survey. This difference may be due in part to the fact that the commercial catch sampled came from only five statistical areas and does not include waters closed to commercial fishing. The trawl survey included closed waters where smaller, immature male crab are more likely to be concentrated. Another reason for this difference may be commercial crab pots are not able to catch the smaller, immature crab a trawl will.

COMMENTS ON OBSERVER PROGRAM

This was the third season that independent observers were placed on board crab vessels during the summer commercial fishery in Norton Sound. All five independent observers had prior experience. Three of the five collected all required data. One observer turned in partial data at debriefing because his briefcase fell into the water. An audio cassette containing most of his data was damaged. One other observer did not collect all data due to the small number of crab his processor caught. Bycatch information was considered optional, and four of the six observers collected usable data.

Observers were given a half day orientation on July 30 to familiarize them with the information they were required to collect. This contributed to the high quality of data that was collected through the duration of the fishery. One change that might be made would be on the "Red King Crab Catcher Log Summary". A column should be included for longitude and latitude of each string observed. This would be valuable in determining the distribution of crab in specific statistical areas.

SUMMARY

This years fishing effort was concentrated in statistical area 656401 and 666401. The greatest percent of legal crab harvested was in area 656401. Mean carapace length of harvested legal crab was 199.7 mm and mean weight was 2.9 pounds. This season's legal male new shell/old shell ratio was 71% new shell to 29% old shell. Post recruit crab dominated the harvest - 71.5% of the harvested stock. Recruits composed 28.5% of the harvest. This was an increase in recruitment from 1990. The percent of female crab with a high degree of ovigerity was 73%. This was a decline from the previous years. Female ovigerity has undergone a moderate but steady decline during the period from 1987 to 1992 - 92% of adult female crab having a high degree of ovigerity in 1987 and 73% in 1992. Observers also noted most eggs were dark purple in color. Illegal harvest was 0.08%, well below the 3% guideline thought to be citable by Fish and Wildlife Protection. The onboard observer program is thought to contribute to this low level of illegal harvest.

LITERATURE CITED

- Alaska Department of Fish and Game. 1989. Alaska Department of Fish and Game observer manual for Alaskan crab processors. Alaska Department of Fish and Game, Division of Commercial Fisheries, 211 Mission Road, Kodiak, Alaska.
- Bue, F.J. and C.F. Lean. 1989. Norton Sound winter red king crab studies, 1989. Regional Information Report No 3N90-05. Alaska Department of Fish and Game, Division of Commercial Fisheries, Nome, Alaska. 22pp.
- Gebhard J.G. 1990. Norton Sound summer commercial red king crab fishery observer project report, 1990. Regional Information Report No. 3N90-23. Alaska Department of Fish and Game, Division of Commercial Fisheries, Nome, Alaska. 40pp.
- Gebhard J.G. and C.F. Lean. 1989. Norton Sound summer commercial red king crab fishery observer project report, 1989. Regional Information Report No. 3N89-26. Alaska Department of Fish and Game, Division of Commercial Fisheries, Nome, Alaska. 37pp.
- Gebhard J.G. and C.F. Lean. 1988. Norton Sound summer commercial red king crab fishery observer project report, 1988. Regional Information Report No. 3N88-32. Alaska Department of Fish and Game, Division of Commercial Fisheries, Nome, Alaska. 45pp.
- Knuepfer G.R. and J.G. Gebhard. 1990. Norton Sound winter red king crab studies, 1990. Regional Information Report No. 3N90-19. Alaska Department of Fish and Game, Division of Commercial Fisheries, Nome, Alaska. 25pp.
- Lean, C.F. 1987. Catch rates, size composition and growth of red king crab taken in Norton Sound near Nome during the winter of 1987. AYK Region Shellfish Report #12. Alaska Department of Fish and Game, Division of Commercial Fisheries, Nome, Alaska. 13pp.
- Sandone, G.J., F.J. Bue and S.E. Merkouris. 1988. Norton Sound summer commercial red king crab fishery observer project report, 1987. Regional Information Report No. 3N88-08. Alaska Department of Fish and Game, Division of Commercial Fisheries, Nome, Alaska. 26pp.

Table 1. Red king crab verbal catch reports for all fishing vessels by statistical area,
Norton Sound Section, Eastern Bering Sea, for August 1, 12p.m. through
August 3, 8 a.m. 1992.

Date	656401		666401		636401		656330		646330		646402	
	# pots	# crab	# pots	# crab	# pots	# crab	# pots	# crab	# pots	# crab	# pots	# crab
8/1-8/3	1550	6722	1696	3340	558	846	599	3248	25	692	118	1127
Catch per pot (CPP)		4.3		2.0		1.5		5.4		27.7		9.6
Percent of harvest (%)		42.1		20.9		5.3		20.3		4.3		7.1
Percent of pots pulled (%)		34.1		37.3		12.3		13.2		0.5		2.6
Total Pots		4546										
Total Harvest		15975										
CPP		3.5										

Table 2. Red king crab season total catch (from fish ticket reports) by statistical area for Norton Sound Section, Eastern Bering Sea, August 1 – 3, 1992.

Statistical Area	Number	Pounds	Pots	Catch per pot (CPP)	Average Weight (Lbs.)	Percent pots pulled in stat. area (%)	Percent harvest in stat. area (%)
656401	18193	53119	2714	6.7	2.92	47.2	73.1
666401	3351	10632	1804	1.9	3.17	31.4	13.5
636401	368	1159	419	0.9	3.15	7.3	1.5
656330	1646	4814	421	3.9	2.92	7.3	6.6
666330	1344	4305	388	3.5	3.2	6.8	5.4
Total	24902	74029	5746	4.3	3.0	100	100

Table 3. Observer crab catch summary; number of observed pot lifts, average soak time, number of legal males, sublegal males, and females and proportion (%) of each captured by statistical area, and the corresponding mean number of crab caught per observed pot lift (CPP) and pot lift standardized to a 24 hour soak period (CPP/24hr), Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Stat. Area	#Pot Lifts Observed	Ave. Soak Time	Legal Males				Sublegal males				Females			
			number	%	CPP	CPP /24h	number	%	CPP	CPP /24h	number	%	CPP	CPP /24h
656401	211	14.5	426	75.4%	2.0	3.3	124	21.9%	0.6	1.0	15	2.7%	0.1	0.1
666401	381	13.9	854	71.0%	2.2	3.9	254	21.1%	0.7	1.2	95	7.9%	0.2	0.4
636401	102	19.8	59	34.7%	0.6	0.7	74	43.5%	0.7	0.9	37	21.8%	0.4	0.4
656330	39	20.8	81	92.0%	2.1	2.4	7	8.0%	0.2	0.2	0	0.0%	0.0	0.0
TOTAL	733	15.3	1420		1.9	3.0	459		0.6	1.0	147		0.2	0.3
Percent of obs. catch				70.1%				22.7%				7.3%		
Total crab obs.			2026											

Table 4. Carapace length measurement summary of sampled legal male red king crab captured during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992

Carapace Length (mm)	New shell			Old shell			Total		
	No.	Ave Length Calc.	%	No.	Ave Length Calc.	%	No.	Ave Length Calc.	%
98		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
99	1	0.05	0.1%	1	0.13	0.1%	2	0.08	0.1%
100	3	0.16	0.2%		0.00	0.0%	3	0.12	0.1%
101	8	0.33	0.3%		0.00	0.0%	6	0.24	0.2%
102	12	0.67	0.7%	2	0.27	0.3%	14	0.56	0.5%
103	22	1.25	1.2%	4	0.55	0.5%	26	1.04	1.0%
104	20	1.14	1.1%	9	1.25	1.2%	29	1.18	1.1%
105	42	2.42	2.3%	17	2.39	2.3%	59	2.41	2.3%
106	39	2.27	2.1%	10	1.42	1.3%	49	2.02	1.9%
107	54	3.17	3.0%	10	1.43	1.3%	64	2.67	2.5%
108	53	3.15	2.9%	19	2.75	2.5%	72	3.03	2.8%
109	51	3.05	2.8%	18	2.83	2.4%	69	2.93	2.7%
110	63	3.81	3.5%	26	3.83	3.5%	89	3.82	3.5%
111	57	3.48	3.1%	28	4.17	3.8%	85	3.68	3.3%
112	75	4.62	4.1%	37	5.55	5.0%	112	4.89	4.4%
113	81	3.79	3.4%	29	4.39	3.9%	90	3.95	3.5%
114	75	4.70	4.1%	38	5.81	5.1%	113	5.02	4.4%
115	96	6.07	5.3%	48	7.08	6.2%	142	6.36	5.5%
116	55	3.51	3.0%	31	4.82	4.2%	86	3.89	3.4%
117	55	3.54	3.0%	27	4.23	3.8%	82	3.74	3.2%
118	81	5.25	4.5%	39	6.17	5.2%	120	5.52	4.7%
119	55	3.60	3.0%	25	3.99	3.4%	80	3.71	3.1%
120	76	5.01	4.2%	43	6.92	5.8%	119	5.57	4.8%
121	71	4.72	3.9%	23	3.73	3.1%	94	4.43	3.7%
122	66	4.42	3.6%	35	5.72	4.7%	101	4.80	3.9%
123	52	3.51	2.9%	24	3.96	3.2%	76	3.84	3.0%
124	48	3.27	2.6%	19	3.16	2.5%	67	3.24	2.6%
125	49	3.37	2.7%	22	3.69	2.9%	71	3.46	2.8%
126	38	2.63	2.1%	17	2.67	2.3%	55	2.70	2.1%
127	50	3.49	2.7%	10	1.70	1.3%	60	2.97	2.3%
128	26	1.83	1.4%	19	3.26	2.5%	45	2.24	1.8%
129	34	2.41	1.9%	13	2.25	1.7%	47	2.38	1.8%
130	24	1.71	1.3%	18	2.79	2.1%	40	2.09	1.6%
131	31	2.23	1.7%	14	2.48	1.9%	45	2.30	1.8%
132	32	2.32	1.8%	10	1.77	1.3%	42	2.16	1.6%
133	18	1.32	1.0%	13	2.32	1.7%	31	1.81	1.2%
134	19	1.40	1.0%	9	1.82	1.2%	28	1.48	1.1%
135	21	1.56	1.2%	5	0.90	0.7%	26	1.37	1.0%
136	19	1.42	1.0%	9	1.64	1.2%	28	1.48	1.1%
137	20	1.51	1.1%	1	0.16	0.1%	21	1.12	0.8%
138	19	1.44	1.0%		0.00	0.0%	19	1.02	0.7%
139	12	0.92	0.7%	3	0.56	0.4%	15	0.81	0.6%
140	18	1.38	1.0%	5	0.94	0.7%	23	1.25	0.9%
141	16	1.24	0.9%	1	0.19	0.1%	17	0.93	0.7%
142	16	1.25	0.9%	1	0.19	0.1%	17	0.94	0.7%
143	11	0.86	0.6%	1	0.19	0.1%	12	0.87	0.5%
144	9	0.71	0.5%	2	0.39	0.3%	11	0.62	0.4%
145	12	0.96	0.7%	4	0.78	0.5%	16	0.90	0.6%
146	4	0.32	0.2%		0.00	0.0%	4	0.23	0.2%
147	5	0.40	0.3%	1	0.20	0.1%	6	0.34	0.2%
148	7	0.57	0.4%	5	0.99	0.7%	12	0.69	0.5%
149	4	0.33	0.2%	1	0.20	0.1%	5	0.29	0.2%
150	2	0.16	0.1%		0.00	0.0%	2	0.12	0.1%
151	4	0.33	0.2%	2	0.40	0.3%	6	0.35	0.2%
152	2	0.17	0.1%	1	0.20	0.1%	3	0.18	0.1%
153	1	0.08	0.1%		0.00	0.0%	1	0.08	0.0%
154	2	0.17	0.1%	1	0.21	0.1%	3	0.18	0.1%
155	1	0.09	0.1%		0.00	0.0%	1	0.06	0.0%
156		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
157		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
158	1	0.09	0.1%		0.00	0.0%	1	0.06	0.0%
159		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
160		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
161		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
162		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
163		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
164	2	0.18	0.1%		0.00	0.0%	2	0.13	0.1%
165	1	0.09	0.1%		0.00	0.0%	1	0.06	0.0%
170	1	0.09	0.1%		0.00	0.0%	1	0.07	0.0%
171		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
Total No.	1820		70.9%	748		29.1%	2568		100.0%
Mean		120.0			119.2			119.7	
Total legal			2566						
Total Recruits			730						
Percent			28.4%						
Total Post Recruits			1836						
Percent			71.6%						

Table 5. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 656401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3 1992.

Carapace Length (mm)	New shell			Old shell			Total		
	No.	Ave Length Calc.	%	No.	Ave Length Calc.	%	No.	Ave Length Calc.	%
96		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
99		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
100	2	0.35	0.4%		0.00	0.0%	2	0.23	0.2%
101	2	0.36	0.4%		0.00	0.0%	2	0.24	0.2%
102	7	1.28	1.2%		0.00	0.0%	7	0.84	0.8%
103	5	0.91	0.9%	2	0.72	0.7%	7	0.85	0.8%
104	8	1.10	1.1%	7	2.53	2.4%	13	1.58	1.5%
105	17	3.16	3.0%	5	1.82	1.7%	22	2.71	2.6%
106	14	2.63	2.5%	5	1.84	1.7%	19	2.36	2.2%
107	27	5.11	4.8%	4	1.49	1.4%	31	3.89	3.8%
108	17	3.25	3.0%	5	1.68	1.7%	22	2.79	2.6%
109	21	4.05	3.7%	8	2.27	2.1%	27	3.45	3.2%
110	26	6.06	4.6%	10	3.82	3.5%	36	4.64	4.2%
111	18	3.54	3.2%	10	3.85	3.5%	28	3.64	3.3%
112	26	5.15	4.8%	21	8.17	7.3%	47	6.17	5.5%
113	26	5.20	4.8%	12	4.71	4.2%	38	5.03	4.5%
114	21	4.24	3.7%	19	7.52	6.8%	40	5.35	4.7%
115	31	6.31	6.5%	24	9.58	8.3%	55	7.42	6.4%
116	17	3.46	3.0%	15	8.04	5.2%	32	4.35	3.8%
117	25	6.10	4.4%	10	4.08	3.5%	35	4.80	4.1%
118	24	5.01	4.2%	13	5.83	4.5%	37	5.12	4.3%
119	25	5.27	4.4%	13	5.37	4.5%	38	5.30	4.5%
120	26	5.52	4.6%	18	7.50	6.3%	44	6.19	5.2%
121	22	4.71	3.9%	5	2.10	1.7%	27	3.83	3.2%
122	21	4.53	3.7%	18	7.83	6.3%	39	5.58	4.8%
123	14	3.05	2.5%	10	4.27	3.5%	24	3.46	2.8%
124	11	2.41	1.9%	9	3.88	3.1%	20	2.91	2.3%
125	10	3.54	2.8%	8	3.47	2.8%	24	3.52	2.8%
126	11	2.45	1.9%	4	1.75	1.4%	15	2.22	1.8%
127	11	2.47	1.9%	5	2.20	1.7%	16	2.38	1.9%
128	5	1.13	0.9%	6	2.67	2.1%	11	1.65	1.3%
129	10	2.28	1.8%	1	0.45	0.3%	11	1.66	1.3%
130	4	0.92	0.7%	2	0.90	0.7%	6	0.91	0.7%
131	6	1.39	1.1%	3	1.36	1.0%	9	1.38	1.1%
132	7	1.64	1.2%	4	1.83	1.4%	11	1.70	1.3%
133	2	0.47	0.4%	2	0.92	0.7%	4	0.62	0.5%
134	2	0.47	0.4%	6	2.79	2.1%	8	1.26	0.9%
135	4	0.98	0.7%		0.00	0.0%	4	0.63	0.5%
136	5	1.20	0.9%	3	1.42	1.0%	8	1.28	0.9%
137	4	0.97	0.7%		0.00	0.0%	4	0.64	0.5%
138	5	1.22	0.9%		0.00	0.0%	5	0.81	0.6%
139	1	0.25	0.2%		0.00	0.0%	1	0.18	0.1%
140	4	0.99	0.7%	1	0.49	0.3%	5	0.82	0.6%
141	3	0.75	0.5%		0.00	0.0%	3	0.50	0.4%
142	6	2.01	1.4%		0.00	0.0%	6	1.33	0.9%
143	1	0.25	0.2%		0.00	0.0%	1	0.17	0.1%
144	1	0.25	0.2%		0.00	0.0%	1	0.17	0.1%
145	1	0.26	0.2%	1	0.50	0.3%	2	0.34	0.2%
146		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
147		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
148	1	0.26	0.2%		0.00	0.0%	1	0.17	0.1%
149	1	0.26	0.2%		0.00	0.0%	1	0.17	0.1%
150		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
151		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
152	1	0.27	0.2%		0.00	0.0%	1	0.18	0.1%
153		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
154		0.00	0.0%	1	0.53	0.3%	1	0.18	0.1%
155		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
156		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
157		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
158		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
159		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
160		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
161		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
162		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
163		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
164		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
165		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
Total No.	565		86.2%	288		33.8%	853		100.0%
Mean		117.5			117.6			117.5	
Total legal			653						
Total Recruits			266						
Percent			31.2%						
Total Post Recruits			587						
Percent			68.8%						

Table 6. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 566401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3 1992.

Carapace Length (mm)	New shell			Old shell			Total	
	No.	Ave Length Calc.	%	No.	Ave Length Calc.	%	No.	Ave Length Calc. %
98		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
99	1	0.22	0.2%		0.00	0.0%	1	0.17 0.2%
100	1	0.22	0.2%		0.00	0.0%	1	0.17 0.2%
101	1	0.22	0.2%		0.00	0.0%	1	0.17 0.2%
102	1	0.22	0.2%	2	1.40	1.4%	3	0.51 0.5%
103	9	2.04	2.0%	3	2.12	2.1%	12	2.08 2.0%
104	5	1.15	1.1%	1	0.71	0.7%	6	1.04 1.0%
105	8	1.85	1.8%	6	4.32	4.1%	14	2.45 2.3%
106	12	2.80	2.6%	2	1.45	1.4%	14	2.47 2.3%
107	10	2.36	2.2%	2	1.47	1.4%	12	2.14 2.0%
108	12	2.85	2.6%	4	2.96	2.7%	16	2.86 2.7%
109	9	2.18	2.0%	1	0.75	0.7%	10	1.82 1.7%
110	11	2.67	2.4%	5	3.77	3.4%	16	2.93 2.7%
111	17	4.16	3.7%	4	3.04	2.7%	21	3.89 3.5%
112	25	6.17	5.5%	3	2.90	2.1%	28	5.23 4.7%
113	6	1.99	1.8%	4	3.10	2.7%	12	2.28 2.0%
114	25	6.26	5.5%	5	3.90	3.4%	30	5.70 5.0%
115	23	6.03	5.1%	7	5.61	4.8%	30	5.75 5.0%
116	13	3.32	2.9%	6	8.38	5.5%	21	4.08 3.5%
117	8	2.06	1.8%	7	6.81	4.8%	15	2.93 2.5%
118	17	4.42	3.7%	12	9.70	8.2%	29	5.70 4.8%
119	12	3.15	2.8%	4	3.26	2.7%	16	3.17 2.7%
120	18	4.78	4.0%	3	2.47	2.1%	21	4.20 3.5%
121	14	3.73	3.1%	6	4.97	4.1%	20	4.03 3.3%
122	13	3.49	2.9%	5	4.18	3.4%	18	3.68 3.0%
123	13	3.52	2.9%	5	4.21	3.4%	18	3.69 3.0%
124	15	4.10	3.3%	4	3.40	2.7%	19	3.93 3.2%
125	9	2.48	2.0%	6	5.14	4.1%	15	3.13 2.5%
126	6	2.22	1.8%	4	3.45	2.7%	12	2.52 2.0%
127	16	4.48	3.5%	2	1.74	1.4%	18	3.61 3.0%
128	9	2.54	2.0%	4	3.51	2.7%	13	2.77 2.2%
129	10	2.64	2.2%	5	4.42	3.4%	15	3.23 2.5%
130	9	2.58	2.0%	4	3.58	2.7%	13	2.82 2.2%
131	4	1.15	0.9%	2	1.79	1.4%	6	1.31 1.0%
132	8	2.33	1.8%	1	0.90	0.7%	9	1.98 1.5%
133	8	2.34	1.8%	2	1.82	1.4%	10	2.22 1.7%
134	4	1.18	0.9%	2	1.84	1.4%	6	1.34 1.0%
135	6	1.78	1.3%	3	2.77	2.1%	9	2.03 1.5%
136	5	1.50	1.1%		0.00	0.0%	5	1.13 0.8%
137	5	1.51	1.1%	1	0.94	0.7%	6	1.37 1.0%
138	4	1.22	0.9%		0.00	0.0%	4	0.92 0.7%
139	5	1.53	1.1%	2	1.90	1.4%	7	1.62 1.2%
140	7	2.18	1.5%	3	2.88	2.1%	10	2.33 1.7%
141	9	2.60	2.0%		0.00	0.0%	9	2.12 1.5%
142	4	1.25	0.9%	1	0.97	0.7%	5	1.18 0.8%
143	4	1.26	0.9%		0.00	0.0%	4	0.95 0.7%
144	2	0.63	0.4%		0.00	0.0%	2	0.46 0.3%
145	5	1.60	1.1%		0.00	0.0%	5	1.21 0.8%
146	1	0.32	0.2%		0.00	0.0%	1	0.24 0.2%
147	2	0.65	0.4%		0.00	0.0%	2	0.49 0.3%
148	3	0.96	0.7%	1	1.01	0.7%	4	0.99 0.7%
149		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
150	1	0.33	0.2%		0.00	0.0%	1	0.25 0.2%
151		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
152		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
153		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
154	1	0.34	0.2%		0.00	0.0%	1	0.26 0.2%
155	1	0.34	0.2%		0.00	0.0%	1	0.26 0.2%
156		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
157		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
158	1	0.35	0.2%		0.00	0.0%	1	0.26 0.2%
159		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
160		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
161		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
162		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
163		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
164		0.00	0.0%		0.00	0.0%	0	0.00 0.0%
165	1	0.36	0.2%		0.00	0.0%	1	0.28 0.2%
170	1	0.37	0.2%		0.00	0.0%	1	0.28 0.2%
Total No.	454		75.7%	146		24.3%	600	100.0%
Mean		121.1			119.5			120.7
Total legal			600					
Total Recruits Percent			178 29.7%					
Total Post Recruits Percent			422 70.3%					

Table 7. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 638401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3 1992.

Carapace Length (mm)	New shell			Old shell			Total		
	No.	Ave Length	%	No.	Ave Length	%	No.	Ave Length	%
98		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
99		0.00	0.0%	1	2.58	2.7%	1	0.93	0.9%
100		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
101	1	1.40	1.4%		0.00	0.0%	1	0.95	0.9%
102		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
103	1	1.49	1.4%		0.00	0.0%	1	0.97	0.9%
104	1	1.51	1.4%		0.00	0.0%	1	0.98	0.9%
105	1	1.52	1.4%	2	5.68	5.4%	3	2.97	2.8%
106	1	1.54	1.4%		0.00	0.0%	1	1.00	0.9%
107	1	1.55	1.4%		0.00	0.0%	1	1.01	0.9%
108		0.00	0.0%	1	2.92	2.7%	1	1.02	0.9%
109	2	3.18	2.9%		0.00	0.0%	2	2.06	1.9%
110	1	1.59	1.4%	1	2.97	2.7%	2	2.08	1.9%
111	2	3.22	2.9%		0.00	0.0%	2	2.09	1.9%
112	5	6.12	7.2%		0.00	0.0%	5	5.28	4.7%
113	3	4.91	4.3%	2	6.11	5.4%	5	5.33	4.7%
114	2	3.30	2.9%	1	3.08	2.7%	3	3.23	2.8%
115	4	6.67	5.8%	1	3.11	2.7%	5	6.42	4.7%
116	4	6.72	5.8%		0.00	0.0%	4	4.38	3.8%
117	2	3.39	2.9%		0.00	0.0%	2	2.21	1.9%
118	5	8.55	7.2%	1	3.19	2.7%	6	6.88	5.7%
119	4	6.90	5.8%	5	18.08	13.5%	9	10.10	8.5%
120	1	1.74	1.4%	2	6.49	5.4%	3	3.40	2.8%
121	2	3.51	2.9%	1	3.27	2.7%	3	3.42	2.8%
122	2	3.54	2.9%	3	9.69	8.1%	5	5.75	4.7%
123	4	7.13	5.8%	1	3.32	2.7%	5	5.80	4.7%
124	1	1.80	1.4%		0.00	0.0%	1	1.17	0.9%
125	3	5.43	4.3%		0.00	0.0%	3	3.54	2.8%
126	2	3.65	2.9%	1	3.41	2.7%	3	3.57	2.8%
127	1	1.84	1.4%		0.00	0.0%	1	1.20	0.9%
128	2	3.71	2.9%	2	6.92	5.4%	4	4.83	3.8%
129		0.00	0.0%	2	6.97	5.4%	2	2.43	1.9%
130	2	3.77	2.9%	1	3.51	2.7%	3	3.66	2.8%
131	1	1.90	1.4%	2	7.08	5.4%	3	3.71	2.8%
132	1	1.91	1.4%		0.00	0.0%	1	1.25	0.9%
133		0.00	0.0%	1	3.59	2.7%	1	1.25	0.9%
134	2	3.88	2.9%		0.00	0.0%	2	2.53	1.9%
135		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
136		0.00	0.0%	1	3.68	2.7%	1	1.28	0.9%
137	2	3.97	2.9%		0.00	0.0%	2	2.58	1.9%
138		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
139		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
140	1	2.03	1.4%	1	3.76	2.7%	2	2.84	1.9%
141		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
142	1	2.06	1.4%		0.00	0.0%	1	1.34	0.9%
143		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
144	1	2.09	1.4%	1	3.89	2.7%	2	2.72	1.9%
145		0.00	0.0%	1	3.92	2.7%	1	1.37	0.9%
146		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
147		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
148		0.00	0.0%	1	4.00	2.7%	1	1.40	0.9%
149		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
150		0.00	0.0%	1	4.05	2.7%	1	1.42	0.9%
151		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
152		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
153		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
154		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
155		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
156		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
157		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
158		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
159		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
160		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
161		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
162		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
163		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
164		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
165		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
Total No.	89	65.1%		37	34.9%		106	100.0%	
Mean		119.6			123.6			120.9	
Total legal			106						
Total Recruits			25						
Percent			23.6%						
Total Post Recruits			81						
Percent			75.4%						

Table 8. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 656330 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3 1992.

Carapace Length (mm)	New shell			Old shell			Total		
	No.	Ave Length	%	No.	Ave Length	%	No.	Ave Length	%
98		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
99		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
100		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
101		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
102		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
103		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
104	1	0.78	0.7%		0.00	0.0%	1	0.58	0.6%
105	1	0.77	0.7%		0.00	0.0%	1	0.59	0.6%
106	1	0.77	0.7%		0.00	0.0%	1	0.80	0.8%
107	4	3.12	2.9%	1	2.61	2.4%	5	3.01	2.6%
108	1	0.79	0.7%	2	5.27	4.9%	3	1.82	1.7%
109	1	0.80	0.7%	1	2.66	2.4%	2	1.22	1.1%
110	3	2.41	2.2%	1	2.68	2.4%	4	2.47	2.2%
111	3	2.43	2.2%	1	2.71	2.4%	4	2.49	2.2%
112	2	1.64	1.5%		0.00	0.0%	2	1.26	1.1%
113	5	4.12	3.6%		0.00	0.0%	5	3.17	2.6%
114	3	2.50	2.2%	2	5.58	4.9%	5	3.20	2.8%
115	2	1.68	1.5%		0.00	0.0%	2	1.29	1.1%
116	3	2.54	2.2%	2	5.65	4.9%	5	3.26	2.8%
117	3	2.56	2.2%		0.00	0.0%	3	1.97	1.7%
118	3	2.58	2.2%	2	5.76	4.9%	5	3.31	2.8%
119	3	2.61	2.2%		0.00	0.0%	3	2.01	1.7%
120	6	5.26	4.4%	1	2.93	2.4%	7	4.72	3.9%
121	6	5.30	4.4%	1	2.95	2.4%	7	4.76	3.9%
122	5	4.45	3.8%	1	2.98	2.4%	6	4.11	3.4%
123	5	4.49	3.8%	3	9.00	7.3%	8	5.53	4.5%
124	3	2.72	2.2%	2	8.05	4.9%	5	3.48	2.8%
125	4	3.65	2.9%		0.00	0.0%	4	2.81	2.2%
126	3	2.76	2.2%	1	3.07	2.4%	4	2.83	2.2%
127	5	4.64	3.6%		0.00	0.0%	5	3.57	2.8%
128	3	2.80	2.2%	2	8.24	4.9%	5	3.60	2.8%
129	4	3.77	2.9%	1	3.15	2.4%	5	3.62	2.8%
130	1	0.95	0.7%	1	3.17	2.4%	2	1.48	1.1%
131	6	5.74	4.4%	2	6.39	4.9%	8	5.89	4.5%
132	5	4.82	3.8%	1	3.22	2.4%	6	4.45	3.4%
133	1	0.97	0.7%	1	3.24	2.4%	2	1.49	1.1%
134	1	0.98	0.7%		0.00	0.0%	1	0.75	0.6%
135	2	1.97	1.5%	1	3.29	2.4%	3	2.28	1.7%
136	2	1.99	1.5%	1	3.32	2.4%	3	2.29	1.7%
137	2	2.00	1.5%		0.00	0.0%	2	1.54	1.1%
138	2	2.01	1.5%		0.00	0.0%	2	1.55	1.1%
139	2	2.03	1.5%		0.00	0.0%	2	1.56	1.1%
140	2	2.04	1.5%		0.00	0.0%	2	1.57	1.1%
141	2	2.08	1.5%		0.00	0.0%	2	1.58	1.1%
142	2	2.07	1.5%		0.00	0.0%	2	1.60	1.1%
143	2	2.09	1.5%	1	3.49	2.4%	3	2.41	1.7%
144	2	2.10	1.5%	1	3.51	2.4%	3	2.43	1.7%
145	3	3.16	2.2%	1	3.54	2.4%	4	3.26	2.2%
146	2	2.13	1.5%		0.00	0.0%	2	1.64	1.1%
147	2	2.15	1.5%	1	3.59	2.4%	3	2.48	1.7%
148	3	3.24	2.2%	3	10.83	7.3%	6	4.99	3.4%
149	2	2.18	1.5%	1	3.63	2.4%	3	2.51	1.7%
150	1	1.09	0.7%		0.00	0.0%	1	0.84	0.6%
151	3	3.31	2.2%	1	3.65	2.4%	4	3.39	2.2%
152	1	1.11	0.7%	1	3.71	2.4%	2	1.71	1.1%
153	1	1.12	0.7%		0.00	0.0%	1	0.86	0.6%
154	1	1.12	0.7%		0.00	0.0%	1	0.87	0.6%
155		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
156		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
157		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
158		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
159		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
160		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
161		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
162		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
163		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
164	1	1.20	0.7%		0.00	0.0%	1	0.92	0.6%
165		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
Total No.	137		77.0%	41		23.0%	178		100.0%
Mean		127.5			127.8			127.8	
Total legal			178						
Total Recruits			27						
Percent			15.2%						
Total Post Recruits			151						
Percent			84.8%						

Table 9. Carapace length measurement summary of sampled legal male red king crab captured in statistical area 656330 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3 1992.

Carapace Length (mm)	New shell			Old shell			Total		
	No.	Ave Length Calc.	%	No.	Ave Length Calc.	%	No.	Ave Length Calc.	%
98		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
99		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
100		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
101		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
102		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
103		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
104	1	2.48	2.4%		0.00	0.0%	1	1.65	1.6%
105		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
106	1	2.52	2.4%		0.00	0.0%	1	1.68	1.6%
107	1	2.56	2.4%		0.00	0.0%	1	1.70	1.6%
108	3	7.71	7.1%	3	15.43	14.3%	6	10.29	9.5%
109	1	2.60	2.4%		0.00	0.0%	1	1.73	1.6%
110	2	5.24	4.8%	1	5.24	4.8%	3	5.24	4.8%
111	1	2.64	2.4%	1	5.29	4.8%	2	3.62	3.2%
112	1	2.67	2.4%	1	5.33	4.8%	2	3.56	3.2%
113	3	8.07	7.1%		0.00	0.0%	3	5.38	4.8%
114	1	2.71	2.4%	1	5.43	4.8%	2	3.62	3.2%
115	2	5.48	4.8%	1	5.48	4.8%	3	5.48	4.8%
116		0.00	0.0%	1	5.52	4.8%	1	1.84	1.6%
117		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
118	2	5.62	4.8%		0.00	0.0%	2	3.75	3.2%
119		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
120	4	11.43	9.5%		0.00	0.0%	4	7.62	6.3%
121	3	8.64	7.1%	3	17.29	14.3%	6	11.52	9.5%
122	1	2.90	2.4%	1	5.81	4.8%	2	3.87	3.2%
123	2	5.68	4.8%		0.00	0.0%	2	3.90	3.2%
124	2	5.90	4.8%	1	5.90	4.8%	3	5.90	4.8%
125	1	2.98	2.4%	2	11.90	9.5%	3	5.95	4.8%
126	1	3.00	2.4%		0.00	0.0%	1	2.00	1.6%
127	2	8.05	4.8%	1	8.05	4.8%	3	8.05	4.8%
128		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
129		0.00	0.0%	1	6.14	4.8%	1	2.05	1.8%
130	1	3.10	2.4%	1	6.19	4.8%	2	4.13	3.2%
131	3	9.36	7.1%		0.00	0.0%	3	6.24	4.8%
132	1	3.14	2.4%		0.00	0.0%	1	2.10	1.6%
133		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
134	1	3.19	2.4%		0.00	0.0%	1	2.13	1.6%
135		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
136		0.00	0.0%	2	12.95	9.5%	2	4.32	3.2%
137		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
138	1	3.29	2.4%		0.00	0.0%	1	2.19	1.6%
139		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
140		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
141		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
142		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
143		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
144		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
145		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
146		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
147		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
148		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
149		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
150		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
151		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
152		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
153		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
154		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
155		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
156		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
157		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
158		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
159		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
160		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
161		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
162		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
163		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
164		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
165		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
Total No.	42		66.7%	21		33.3%	63		100.0%
Mean		119.1			119.9			119.2	
Total legal			63						
Total Recruits			17						
Percent			27.0%						
Total Post Recruits			46						
Percent			73.0%						

Table 10. Percent of newly recruited male king crab by statistical area,
Norton Sound Section, Eastern Bering Sea, August 1 – 3, 1992.

Statistical Area	# Crab	%Recruits	Range of % Recruits	# of vessel days sampled
656401	853	31%	22–42%	4
666401	600	30%	11–46%	3
636401	106	23%	14–28%	2
656330	178	15%	6–26%	1
666330	63	27%	27%	1
Total	1800 *	28%	6–46%	11

* total number of crab does not equal the same total number of legal crab in Table 4
because some length frequency forms had no statistical area information recorded.

Table 11. Carapace length measurement summary of Pre-recruit, Recruit, and Post-recruit male and female crab, captured during the summer and fall crab harvest, Norton Sound Section, Eastern Bering Sea, August 1 - 3, 1982.

CARAPACE FACE LTH. (mm)	PRE-RECRUIT (Subadults)			RECRUIT			POST-RECRUIT			TOTALS	
	No.	FREQ.	%	No.	FREQ.	%	No.	FREQ.	%	No.	FREQ.
53	2	0.04%	0	0.00%	0	0.00%	0	0.00%	0	2	0.04%
55	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0	0	0.00%
56	1	0.02%	0	0.00%	0	0.00%	0	0.00%	1	0.02%	1
57	1	0.02%	0	0.00%	0	0.00%	0	0.00%	1	0.02%	1
58	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0
59	1	0.02%	0	0.00%	0	0.00%	0	0.00%	1	0.02%	1
60	2	0.04%	0	0.00%	0	0.00%	0	0.00%	2	0.04%	2
61	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0
62	3	0.07%	0	0.00%	0	0.00%	0	0.00%	3	0.07%	3
63	5	0.11%	0	0.00%	0	0.00%	0	0.00%	5	0.11%	5
64	2	0.04%	0	0.00%	0	0.00%	0	0.00%	2	0.04%	2
65	7	0.15%	0	0.00%	0	0.00%	0	0.00%	7	0.15%	7
66	3	0.07%	0	0.00%	0	0.00%	0	0.00%	3	0.07%	3
67	8	0.18%	0	0.00%	0	0.00%	0	0.00%	8	0.18%	8
68	4	0.13%	0	0.00%	0	0.00%	0	0.00%	8	0.13%	8
69	9	0.20%	0	0.00%	0	0.00%	0	0.00%	9	0.20%	9
70	3	0.07%	0	0.00%	0	0.00%	0	0.00%	3	0.07%	3
71	11	0.25%	0	0.00%	0	0.00%	0	0.00%	11	0.25%	11
72	7	0.16%	0	0.00%	0	0.00%	0	0.00%	7	0.16%	7
73	12	0.27%	0	0.00%	0	0.00%	0	0.00%	12	0.27%	12
74	8	0.18%	0	0.00%	0	0.00%	0	0.00%	8	0.18%	8
75	7	0.16%	0	0.00%	0	0.00%	0	0.00%	7	0.16%	7
76	8	0.18%	0	0.00%	0	0.00%	0	0.00%	8	0.18%	8
77	10	0.22%	0	0.00%	0	0.00%	0	0.00%	10	0.22%	10
78	12	0.27%	0	0.00%	0	0.00%	0	0.00%	12	0.27%	12
79	11	0.25%	0	0.00%	0	0.00%	0	0.00%	11	0.25%	11
80	10	0.22%	0	0.00%	0	0.00%	0	0.00%	10	0.22%	10
81	11	0.25%	0	0.00%	0	0.00%	0	0.00%	11	0.25%	11
82	20	0.45%	0	0.00%	0	0.00%	0	0.00%	20	0.45%	20
83	15	0.33%	0	0.00%	0	0.00%	0	0.00%	15	0.33%	15
84	23	0.51%	0	0.00%	0	0.00%	0	0.00%	23	0.51%	23
85	17	0.38%	0	0.00%	0	0.00%	0	0.00%	17	0.38%	17
86	15	0.33%	0	0.00%	0	0.00%	0	0.00%	15	0.33%	15
87	15	0.33%	0	0.00%	0	0.00%	0	0.00%	15	0.33%	15
88	19	0.42%	0	0.00%	0	0.00%	0	0.00%	19	0.42%	19
89	28	0.59%	0	0.00%	0	0.00%	0	0.00%	28	0.59%	28
90	18	0.39%	0	0.00%	0	0.00%	0	0.00%	18	0.39%	18
91	21	0.47%	0	0.00%	0	0.00%	0	0.00%	21	0.47%	21
92	18	0.39%	0	0.00%	0	0.00%	0	0.00%	18	0.39%	18
93	17	0.38%	0	0.00%	0	0.00%	0	0.00%	17	0.38%	17
94	16	0.36%	0	0.00%	0	0.00%	0	0.00%	16	0.36%	16
95	22	0.48%	0	0.00%	0	0.00%	0	0.00%	22	0.48%	22
96	22	0.48%	0	0.00%	0	0.00%	0	0.00%	22	0.48%	22
97	17	0.38%	0	0.00%	0	0.00%	0	0.00%	17	0.38%	17
98	20	0.45%	0	0.00%	0	0.00%	0	0.00%	20	0.45%	20
99	22	0.48%	2	1	0.02%	2	1	0.02%	25	0.56%	25
100	28	0.59%	5	3	0.10%	0	0.00%	31	0.69%	31	0.69%
101	15	0.33%	8	8	0.25%	0	0.00%	24	0.54%	24	0.54%
102	23	0.51%	18	12	0.41%	3	2	0.07%	44	0.96%	44
103	11	0.25%	22	22	0.74%	4	4	0.14%	50	1.12%	50
104	12	0.27%	30	20	0.66%	14	9	0.30%	56	1.25%	56
105	5	0.11%	84	42	1.42%	28	17	0.57%	94	2.10%	94
106	2	0.04%	58	39	1.32%	15	10	0.34%	78	1.70%	78
107	3	0.07%	82	54	1.82%	15	10	0.34%	100	2.23%	100
108	1	0.02%	80	50	1.70%	28	19	0.64%	109	2.43%	109
109	1	0.02%	77	51	1.72%	27	18	0.61%	105	2.35%	105
110	1	0.02%	85	53	1.78%	38	26	0.88%	129	2.87%	129
111	1	0.02%	86	57	1.82%	42	28	0.98%	139	3.07%	139
112	1	0.02%	114	75	2.53%	58	37	1.25%	170	3.78%	170
113	1	0.02%	92	61	2.08%	44	29	0.98%	136	3.04%	136
114	1	0.02%	114	75	2.53%	58	38	1.28%	171	3.82%	171
115	1	0.02%	145	98	3.24%	70	46	1.56%	215	4.80%	215
116	1	0.02%	0	0.00%	130	86	2.90%	130	2.90%	130	2.90%
117	1	0.02%	0	0.00%	124	82	2.77%	124	2.77%	124	2.77%
118	1	0.02%	0	0.00%	182	120	4.05%	182	4.05%	182	4.05%
119	1	0.02%	0	0.00%	121	80	2.70%	121	2.70%	121	2.70%
120	1	0.02%	0	0.00%	180	118	4.02%	180	4.02%	180	4.02%
121	1	0.02%	0	0.00%	142	94	3.17%	142	3.17%	142	3.17%
122	1	0.02%	0	0.00%	153	101	3.41%	153	3.41%	153	3.41%
123	1	0.02%	0	0.00%	115	78	2.57%	115	2.57%	115	2.57%
124	1	0.02%	0	0.00%	101	67	2.28%	101	2.28%	101	2.28%
125	1	0.02%	0	0.00%	108	71	2.42%	108	2.42%	108	2.42%
126	1	0.02%	0	0.00%	80	55	1.88%	80	1.88%	80	1.88%
127	1	0.02%	0	0.00%	91	60	2.03%	91	2.03%	91	2.03%
128	1	0.02%	0	0.00%	88	45	1.52%	88	1.52%	88	1.52%
129	1	0.02%	0	0.00%	71	47	1.59%	71	1.59%	71	1.59%
130	1	0.02%	0	0.00%	61	40	1.35%	61	1.35%	61	1.35%
131	1	0.02%	0	0.00%	88	45	1.52%	88	1.52%	88	1.52%
132	1	0.02%	0	0.00%	84	42	1.42%	84	1.42%	84	1.42%
133	1	0.02%	0	0.00%	47	31	1.03%	47	1.03%	47	1.03%
134	1	0.02%	0	0.00%	42	28	0.95%	42	0.95%	42	0.95%
135	1	0.02%	0	0.00%	39	26	0.89%	39	0.89%	39	0.89%
136	1	0.02%	0	0.00%	42	28	0.95%	42	0.95%	42	0.95%
137	1	0.02%	0	0.00%	32	21	0.71%	32	0.71%	32	0.71%
138	1	0.02%	0	0.00%	29	19	0.64%	29	0.64%	29	0.64%
139	1	0.02%	0	0.00%	23	15	0.51%	23	0.51%	23	0.51%
140	1	0.02%	0	0.00%	35	23	0.78%	35	0.78%	35	0.78%
141	1	0.02%	0	0.00%	28	17	0.57%	28	0.57%	28	0.57%
142	1	0.02%	0	0.00%	26	17	0.57%	26	0.57%	26	0.57%
143	1	0.02%	0	0.00%	18	12	0.41%	18	0.41%	18	0.41%
144	1	0.02%	0	0.00%	17	11	0.37%	17	0.37%	17	0.37%
145	1	0.02%	0	0.00%	24	16	0.54%	24	0.54%	24	0.54%
146	1	0.02%	0	0.00%	4	4	0.14%	4	0.14%	4	0.14%
147	1	0.02%	0	0.00%	8	8	0.27%	8	0.27%	8	0.27%
148	1	0.02%	0	0.00%	18	12	0.41%	18	0.41%	18	0.41%
149	1	0.02%	0	0.00%	8	5	0.17%	8	0.17%	8	0.17%
150	1	0.02%	0	0.00%	3	2	0.07%	3	0.07%	3	0.07%
151	1	0.02%	0	0.00%	9	8	0.27%	9	0.27%	9	0.27%
152	1	0.02%	0	0.00%	5	3	0.10%	5	0.10%	5	0.10%
153	1	0.02%	0	0.00%	2	1	0.03%	2	0.03%	2	0.03%
154	1	0.02%	0	0.00%	5	3	0.10%	5	0.10%	5	0.10%
155	1	0.02%	0	0.00%	2	1	0.03%	2	0.03%	2	0.03%
156	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0
157	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0
158	1	0.02%	0	0.00%	2	1	0.03%	2	0.03%	2	0.03%
159	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0
160	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0
161	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0
162	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0
163	1	0.02%	0	0.00%	0	0.00%	0	0.00%	0	0.00%	0
164	1	0.02%	0	0.00%	3	2	0.07%	3	0.07%	3	0.07%
165	1	0.02%	0	0.00%	2	1	0.03%	2	0.03%	2	0.03%
170	1	0.02%	0	0.00%	2	1	0.03%	2	0.03%	2	0.03%
Total	568	13.34%	1105	730	24.62%	2780	1806	62.01%	4484	100.00%	

Table 12. Observer summary table of legal red king crab weight samples by statistical area, Norton Sound Section, Eastern Bering Sea, August 1–3, 1992.

Statistical Area	# Crab	Weight lbs.	Average Weight (lbs.)	Range of Ave. wts. (lbs.)	# vessel days*
656401	178	542.2	3.1	2.8–3.1	3
666401	155	465.2	3.0	2.4–3.7	3
636401	100	290.5	2.9	2.6–3.2	3
656330	30	88.5	2.9	2.9	1
Total	463	1386.4	3.0	2.4–3.7	10

* Total number of days sampled for weights by observer vessels.

Table 13. Carapace length measurement summary of sampled sublegal male red king crab captured during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Carapace Length (mm)	New Shell			Old Shell			Total		
	No.	Ave Length Calc	%	No.	Ave Length Calc	%	No.	Ave Length Calc	%
52	1	0.10	0.2%	1	0.93	1.8%	2	0.17	0.3%
56	1	0.10	0.2%		0.00	0.0%	1	0.09	0.2%
57	1	0.11	0.2%		0.00	0.0%	1	0.10	0.2%
58		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
59	1	0.11	0.2%		0.00	0.0%	1	0.10	0.2%
60	2	0.22	0.4%		0.00	0.0%	2	0.26	0.3%
61		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
62	3	0.34	0.6%		0.00	0.0%	3	0.31	0.5%
63	5	0.58	0.9%		0.00	0.0%	5	0.53	0.8%
64	2	0.24	0.4%		0.00	0.0%	2	0.21	0.3%
65	7	0.84	1.3%		0.00	0.0%	7	0.76	1.2%
66	3	0.37	0.6%		0.00	0.0%	3	0.33	0.5%
67	8	0.99	1.5%		0.00	0.0%	8	0.90	1.3%
68	6	0.75	1.1%		0.00	0.0%	6	0.68	1.0%
69	8	1.02	1.5%	1	1.23	1.8%	9	1.04	1.5%
70	3	0.39	0.6%		0.00	0.0%	3	0.35	0.5%
71	11	1.44	2.0%		0.00	0.0%	11	1.31	1.8%
72	5	0.66	0.9%	2	2.57	3.6%	7	0.84	1.2%
73	12	1.62	2.2%		0.00	0.0%	12	1.46	2.0%
74	7	0.96	1.3%	1	1.32	1.8%	8	0.99	1.3%
75	5	0.69	0.9%	2	2.68	3.6%	7	0.88	1.2%
76	6	0.84	1.1%	2	2.71	3.6%	8	1.02	1.3%
77	10	1.42	1.8%		0.00	0.0%	10	1.29	1.7%
78	11	1.56	2.0%	1	1.39	1.8%	12	1.57	2.0%
79	9	1.31	1.7%	2	2.82	3.6%	11	1.45	1.8%
80	10	1.48	1.8%		0.00	0.0%	10	1.34	1.7%
81	11	1.64	2.0%		0.00	0.0%	11	1.49	1.8%
82	18	2.72	3.3%	2	2.93	3.6%	20	2.74	3.3%
83	12	1.84	2.2%	3	4.45	5.4%	15	2.08	2.5%
84	22	3.41	4.1%	1	1.50	1.8%	23	3.23	3.8%
85	15	2.35	2.8%	2	3.04	3.6%	17	2.42	2.8%
86	15	2.38	2.8%		0.00	0.0%	15	2.16	2.5%
87	14	2.25	2.6%	1	1.55	1.8%	15	2.18	2.5%
88	16	2.92	3.3%	1	1.57	1.8%	19	2.80	3.2%
89	24	3.94	4.4%	2	3.18	3.6%	26	3.87	4.3%
90	15	2.49	2.8%	1	1.61	1.8%	16	2.41	2.7%
91	19	3.19	3.5%	2	3.25	3.6%	21	3.20	3.5%
92	16	2.72	3.0%		0.00	0.0%	16	2.46	2.7%
93	16	2.75	3.0%	1	1.66	1.8%	17	2.64	2.8%
94	16	2.77	3.0%		0.00	0.0%	16	2.52	2.7%
95	21	3.68	3.9%	1	1.70	1.8%	22	3.49	3.7%
96	20	3.54	3.7%	2	3.43	3.6%	22	3.53	3.7%
97	14	2.51	2.6%	3	5.20	5.4%	17	2.76	2.8%
98	18	3.25	3.3%	2	3.50	3.6%	20	3.28	3.3%
99	19	3.47	3.5%	3	5.30	5.4%	22	3.64	3.7%
100	22	4.06	4.1%	4	7.14	7.1%	26	4.35	4.3%
101	14	2.61	2.6%	1	1.80	1.8%	15	2.53	2.5%
102	18	3.39	3.3%	5	9.11	8.9%	23	3.92	3.8%
103	9	1.71	1.7%	2	3.68	3.6%	11	1.89	1.8%
104	8	1.54	1.5%	4	7.43	7.1%	12	2.09	2.0%
105	5	0.97	0.9%		0.00	0.0%	5	0.88	0.8%
106	2	0.39	0.4%		0.00	0.0%	2	0.35	0.3%
107	2	0.39	0.4%	1	1.91	1.8%	3	0.54	0.5%
108		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
109	1	0.20	0.2%		0.00	0.0%	1	0.18	0.2%
110	1	0.20	0.2%		0.00	0.0%	1	0.18	0.2%
111		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
112		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
113		0.00	0.0%		0.00	0.0%	0	0.00	0.0%
sum	542		90.6%	56		9.4%	598		100.0%
Mean		87.4			90.6			87.7	
Total sublegals		598							

Table 14. Carapace length measurement summary of sampled sublegal male red king crab captured in statistical area 656401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Carapace Length (mm)	New Shell			Old Shell			Total		
	No.	Ave Length Calc	%	No.	Ave Length Calc	%	No.	Ave Length Calc	%
56	1	0.31	0.6%	0	0.00	0.0%	1	0.27	0.5%
57	1	0.31	0.6%	0	0.00	0.0%	1	0.28	0.5%
58	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
59	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
60	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
61	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
62	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
63	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
64	1	0.35	0.6%	0	0.00	0.0%	1	0.31	0.5%
65	1	0.36	0.6%	0	0.00	0.0%	1	0.32	0.5%
66	1	0.36	0.6%	0	0.00	0.0%	1	0.32	0.5%
67	4	1.48	2.2%	0	0.00	0.0%	4	1.31	2.0%
68	1	0.38	0.6%	0	0.00	0.0%	1	0.33	0.5%
69	1	0.38	0.6%	0	0.00	0.0%	1	0.34	0.5%
70	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
71	1	0.39	0.6%	0	0.00	0.0%	1	0.35	0.5%
72	1	0.40	0.6%	0	0.00	0.0%	1	0.35	0.5%
73	1	0.40	0.6%	0	0.00	0.0%	1	0.36	0.5%
74	1	0.41	0.6%	1	3.22	4.3%	2	0.73	1.0%
75	0	0.00	0.0%	1	3.28	4.3%	1	0.37	0.5%
76	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
77	2	0.85	1.1%	0	0.00	0.0%	2	0.75	1.0%
78	8	2.59	3.3%	0	0.00	0.0%	6	2.29	2.9%
79	2	0.87	1.1%	0	0.00	0.0%	2	0.77	1.0%
80	1	0.44	0.6%	0	0.00	0.0%	1	0.39	0.5%
81	2	0.90	1.1%	0	0.00	0.0%	2	0.79	1.0%
82	10	4.53	5.5%	0	0.00	0.0%	10	4.02	4.9%
83	4	1.83	2.2%	0	0.00	0.0%	4	1.63	2.0%
84	4	1.86	2.2%	0	0.00	0.0%	4	1.65	2.0%
85	3	1.41	1.7%	1	3.70	4.3%	4	1.67	2.0%
86	7	3.33	3.9%	0	0.00	0.0%	7	2.95	3.4%
87	3	1.44	1.7%	0	0.00	0.0%	3	1.28	1.5%
88	4	1.94	2.2%	1	3.83	4.3%	5	2.16	2.5%
89	9	4.43	5.0%	1	3.87	4.3%	10	4.36	4.9%
90	7	3.46	3.9%	1	3.91	4.3%	8	3.53	3.9%
91	8	4.02	4.4%	1	3.96	4.3%	9	4.01	4.4%
92	7	3.56	3.9%	0	0.00	0.0%	7	3.16	3.4%
93	10	5.14	5.5%	1	4.04	4.3%	11	5.01	5.4%
94	6	3.12	3.3%	0	0.00	0.0%	6	2.78	2.9%
95	10	5.25	5.5%	1	4.13	4.3%	11	5.12	5.4%
96	8	4.24	4.4%	0	0.00	0.0%	8	3.76	3.9%
97	4	2.14	2.2%	1	4.22	4.3%	5	2.38	2.5%
98	5	2.71	2.8%	0	0.00	0.0%	5	2.40	2.5%
99	9	4.92	5.0%	1	4.30	4.3%	10	4.85	4.9%
100	12	6.63	6.6%	0	0.00	0.0%	12	5.88	5.9%
101	3	1.67	1.7%	1	4.39	4.3%	4	1.98	2.0%
102	6	3.38	3.3%	6	26.61	26.1%	12	6.00	5.9%
103	5	2.85	2.8%	2	8.98	8.7%	7	3.53	3.4%
104	5	2.87	2.8%	3	13.57	13.0%	8	4.08	3.9%
105	2	1.18	1.1%	0	0.00	0.0%	2	1.03	1.0%
106	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
107	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
108	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
109	1	0.60	0.6%	0	0.00	0.0%	1	0.53	0.5%
110	1	0.61	0.6%	0	0.00	0.0%	1	0.54	0.5%
111	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
112	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
113	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
sum	181		88.7%	23		11.3%	204		100.0%
Mean		90.3			96.0			90.9	
Total sublegals		204							

Table 15. Carapace length measurement summary of sampled sublegal male red king crab captured in statistical area 685401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Carapace Length (mm)	New Shell			Old Shell			Total		
	No.	Ave Length Calc	%	No.	Ave Length Calc	%	No.	Ave Length Calc	%
52	1	0.21	0.4%	1	1.93	3.7%	2	0.38	0.7%
57	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
58	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
59	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
60	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
61	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
62	1	0.25	0.4%	0	0.00	0.0%	1	0.23	0.4%
63	1	0.26	0.4%	0	0.00	0.0%	1	0.23	0.4%
64	1	0.26	0.4%	0	0.00	0.0%	1	0.24	0.4%
65	3	0.80	1.2%	0	0.00	0.0%	3	0.72	1.1%
68	1	0.27	0.4%	0	0.00	0.0%	1	0.24	0.4%
67	2	0.55	0.8%	0	0.00	0.0%	2	0.49	0.7%
68	1	0.28	0.4%	1	2.52	3.7%	2	0.50	0.7%
69	3	0.85	1.2%	1	2.56	3.7%	4	1.02	1.5%
70	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
71	2	0.58	0.8%	0	0.00	0.0%	2	0.52	0.7%
72	0	0.00	0.0%	2	5.33	7.4%	2	0.53	0.7%
73	3	0.90	1.2%	0	0.00	0.0%	3	0.81	1.1%
74	1	0.30	0.4%	0	0.00	0.0%	1	0.27	0.4%
75	4	1.23	1.6%	1	2.78	3.7%	5	1.38	1.8%
76	4	1.25	1.6%	2	5.63	7.4%	6	1.68	2.2%
77	6	1.89	2.5%	0	0.00	0.0%	6	1.70	2.2%
78	4	1.28	1.6%	1	2.89	3.7%	5	1.44	1.8%
79	4	1.30	1.6%	2	5.85	7.4%	6	1.75	2.2%
80	6	1.97	2.5%	0	0.00	0.0%	6	1.77	2.2%
81	9	2.99	3.7%	0	0.00	0.0%	9	2.69	3.3%
82	5	1.68	2.0%	2	6.07	7.4%	7	2.12	2.6%
83	7	2.38	2.9%	3	9.22	11.1%	10	3.06	3.7%
84	17	5.85	7.0%	1	3.11	3.7%	18	5.58	6.6%
85	10	3.48	4.1%	1	3.15	3.7%	11	3.45	4.1%
86	7	2.47	2.9%	0	0.00	0.0%	7	2.22	2.6%
87	6	2.14	2.5%	1	3.22	3.7%	7	2.25	2.6%
88	11	3.97	4.5%	0	0.00	0.0%	11	3.57	4.1%
89	14	5.11	5.7%	0	0.00	0.0%	14	4.60	5.2%
90	7	2.58	2.9%	0	0.00	0.0%	7	2.32	2.6%
91	9	3.36	3.7%	0	0.00	0.0%	9	3.02	3.3%
92	8	3.02	3.3%	0	0.00	0.0%	8	2.72	3.0%
93	4	1.52	1.6%	0	0.00	0.0%	4	1.37	1.5%
94	6	2.31	2.5%	0	0.00	0.0%	6	2.08	2.2%
95	8	3.11	3.3%	0	0.00	0.0%	8	2.80	3.0%
98	9	3.54	3.7%	1	3.56	3.7%	10	3.54	3.7%
97	7	2.78	2.9%	2	7.19	7.4%	9	3.22	3.3%
98	11	4.42	4.5%	1	3.63	3.7%	12	4.34	4.4%
99	8	3.25	3.3%	1	3.67	3.7%	9	3.29	3.3%
100	8	3.28	3.3%	3	11.11	11.1%	11	4.06	4.1%
101	8	3.31	3.3%	0	0.00	0.0%	8	2.98	3.0%
102	6	2.51	2.5%	0	0.00	0.0%	6	2.28	2.2%
103	2	0.84	0.8%	0	0.00	0.0%	2	0.76	0.7%
104	3	1.28	1.2%	0	0.00	0.0%	3	1.15	1.1%
105	4	1.72	1.6%	0	0.00	0.0%	4	1.55	1.5%
106	1	0.43	0.4%	0	0.00	0.0%	1	0.39	0.4%
107	1	0.44	0.4%	0	0.00	0.0%	1	0.39	0.4%
108	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
109	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
110	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
111	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
112	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
113	0	0.00	0.0%	0	0.00	0.0%	0	0.00	0.0%
sum	244		90.0%	27		10.0%	271		100.0%
Mean		88.2			83.4			87.7	
Total sublegals		271							

Table 16. Carapace length measurement summary of sampled sublegal male red king crab captured in statistical area 636401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992. (No old shell in sample)

Carapace Length (mm)	New Shell		
	No.	Ave Length Calc	%
57	0	0.00	0.0%
58	0	0.00	0.0%
59	4	3.23	5.5%
60	2	1.64	2.7%
61	0	0.00	0.0%
62	2	1.70	2.7%
63	4	3.45	5.5%
64	0	0.00	0.0%
65	3	2.67	4.1%
66	0	0.00	0.0%
67	2	1.84	2.7%
68	3	2.79	4.1%
69	4	3.78	5.5%
70	3	2.88	4.1%
71	8	7.78	11.0%
72	4	3.95	5.5%
73	8	8.00	11.0%
74	4	4.05	5.5%
75	1	1.03	1.4%
76	2	2.08	2.7%
77	2	2.11	2.7%
78	1	1.07	1.4%
79	2	2.16	2.7%
80	2	2.19	2.7%
81	0	0.00	0.0%
82	2	2.25	2.7%
83	1	1.14	1.4%
84	1	1.15	1.4%
85	1	1.16	1.4%
86	0	0.00	0.0%
87	1	1.19	1.4%
88	1	1.21	1.4%
89	0	0.00	0.0%
90	0	0.00	0.0%
91	0	0.00	0.0%
92	0	0.00	0.0%
93	0	0.00	0.0%
94	1	1.29	1.4%
95	1	1.30	1.4%
96	2	2.63	2.7%
97	0	0.00	0.0%
98	0	0.00	0.0%
99	0	0.00	0.0%
100	0	0.00	0.0%
101	0	0.00	0.0%
102	1	1.40	1.4%
103	0	0.00	0.0%
104	0	0.00	0.0%
105	0	0.00	0.0%
106	0	0.00	0.0%
107	0	0.00	0.0%
108	0	0.00	0.0%
109	0	0.00	0.0%
110	0	0.00	0.0%
111	0	0.00	0.0%
112	0	0.00	0.0%
113	0	0.00	0.0%
sum	73		100.0%
Mean		73.1	
Total sublegals		73	

Table 17. Carapace length measurement summary of sampled sublegal male red king crab captured in statistical area 656330 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992. (No old shell in sample)

Carapace Length (mm)	New Shell		
	No.	Ave Length Calc	%
57	0	0.00	0.0%
58	0	0.00	0.0%
59	0	0.00	0.0%
60	0	0.00	0.0%
61	0	0.00	0.0%
62	0	0.00	0.0%
63	0	0.00	0.0%
64	0	0.00	0.0%
65	0	0.00	0.0%
66	0	0.00	0.0%
67	0	0.00	0.0%
68	0	0.00	0.0%
69	0	0.00	0.0%
70	0	0.00	0.0%
71	0	0.00	0.0%
72	0	0.00	0.0%
73	0	0.00	0.0%
74	0	0.00	0.0%
75	0	0.00	0.0%
76	0	0.00	0.0%
77	0	0.00	0.0%
78	0	0.00	0.0%
79	0	0.00	0.0%
80	0	0.00	0.0%
81	0	0.00	0.0%
82	0	0.00	0.0%
83	0	0.00	0.0%
84	0	0.00	0.0%
85	0	0.00	0.0%
86	0	0.00	0.0%
87	2	24.86	28.6%
88	0	0.00	0.0%
89	0	0.00	0.0%
90	1	12.86	14.3%
91	0	0.00	0.0%
92	0	0.00	0.0%
93	0	0.00	0.0%
94	0	0.00	0.0%
95	0	0.00	0.0%
96	1	13.71	14.3%
97	0	0.00	0.0%
98	0	0.00	0.0%
99	1	14.14	14.3%
100	0	0.00	0.0%
101	0	0.00	0.0%
102	2	29.14	28.6%
103	0	0.00	0.0%
104	0	0.00	0.0%
105	0	0.00	0.0%
106	0	0.00	0.0%
107	0	0.00	0.0%
108	0	0.00	0.0%
109	0	0.00	0.0%
110	0	0.00	0.0%
111	0	0.00	0.0%
112	0	0.00	0.0%
113	0	0.00	0.0%

sum	7	100.0%
Mean	94.7	

Total sublegals 7

Table 18. Carapace length measurement and percent ovigerity summary of sampled female red king crab captured during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Stat.Area	All	Vessel						All	Date						8 1-3 92
JUVENILE	ADULT SIZE PER % OVIGERITY							ADULT SIZE PER % OVIGERITY							
mm no.	mm Full	Hi	Med	Low	0	Sum	mm Full	Hi	Med	Low	0	Sum			
50	60					0	91	1	2			3			
51	61					0	92	1	1			2			
52	62					0	93	1	1	1	1	4			
53	63					0	94	1	1	2		4			
54	64					0	95	3	1	3	1	8			
55	65					0	96	2	4	4		10			
56	66		1			1	97	4	2			6			
57	67			1		1	98					0			
58	68					0	99	1	1			2			
59	69					0	100	1	5			6			
60	70					0	101		3	2		5			
61	71			1		1	102	1	2	1		4			
62	72					0	103		2			2			
63	73			1		1	104		2			2			
64	74	2	1			3	105					0			
65	75		2	1		3	106		2			2			
66	76	1				1	107	1			1	2			
67	77	1	1	1		3	108		2			2			
68	78		1			1	109	1				1			
69	79	1				1	110		1			1			
70	80		1	1		2	111		1			1			
71	81		1			1	112					0			
72	82	1			1	2	113	1				1			
73	83		3		2	5	114		2			2			
74	84	1				1	115		2			2			
75	85	3	2	1	1	7	116		1			1			
76	86	1	3			4	117					0			
77	87	1	1			2	118					0			
78	88			2		2	119					0			
79	89	1	1	2		4	120			2		2			
83	90	2	2	1	1	6	127		1			1			
							128		1			1			
38		15	20	12	5	0	52	19	40	15	3	0	77		
TOTAL ADULTS/OVIGERITY - - -								34	60	27	8	0	129		
% ADULTS/OVIGERITY - - -								26.3	46.5	20.	6.2	0			
MEAN 66.8		WEIGHTED MEAN ADULT LENGTH - - 93.5													
VAR. 30.9		VARIANCE OF THE MEAN LENGTH - - 143.2													

Full=90-100%, Hi=60-89%, Med=30-59%, Lo=1-29%															

Table 19. Carapace length measurement and percent ovigerity summary of sampled female red king crab captured in statistical area 656401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Stat Area 656401		Vessel All						Date 8 1-3 92					

JUVENILE		ADULT SIZE PER % OVIGERITY						ADULT SIZE PER % OVIGERITY					
mm	no.	mm Full	Hi	Med	Low	0	Sum	mm Full	Hi	Med	Low	0	Sum
-----		-----						-----					
50		60					0	91	1				1
51		61					0	92	1				1
52		62					0	93	1	1			2
53		63					0	94			1		1
54		64					0	95					0
55		65					0	96	1		1		2
56		66		1			1	97	2	1			3
57		67					0	98					0
58	1	68					0	99					0
59		69					0	100					0
60		70					0	101					0
61		71					0	102					0
62	1	72					0	103		1			1
63		73					0	104					0
64	3	74					0	105					0
65	1	75					0	106					0
66		76	1				1	107					0
67	1	77					0	108					0
68	1	78					0	109					0
69		79	1				1	110					0
70		80					0	111					0
71		81					0	112					0
72		82	1		1		2	113					0
73	1	83					0	114					0
74		84	1				1	115					0
75	1	85	3				3	116					0
76		86	1				1	117					0
77		87	1				1	118					0
78		88					0	119					0
79		89	1				1	120					0
80		90	1				1	121					0
								122					0
-----		-----						-----					
10		11	1	1	0	0	13	6	3	2	0	0	11
TOTAL ADULTS/OVIGERITY - - -								17	4	3	0	0	24
% ADULTS/OVIGERITY - - - -								70.8	16.6	12.	0	0	
MEAN 66.0		WEIGHTED MEAN ADULT LENGTH -- 88.5											
VAR. 25.3		VARIANCE OF THE MEAN LENGTH -- 66.7											
-----		-----											
Full=90-100%, Hi=60-89%, Med=30-59%, Lo=1-29%													

Table 20. Carapace length measurement and percent ovigerity summary of sampled female red king crab captured in statistical area 666401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Stat Area 666401		Vessel All						Date 8 1~3 92						
JUVENILE		ADULT SIZE PER % OVIGERITY						ADULT SIZE PER % OVIGERITY						
mm	no.	mm Full	Hi	Med	Low	0	Sum	mm Full	Hi	Med	Low	0	Sum	
50		60					0	91	1				1	
51		61					0	92	1				1	
52		62					0	93		1	1		2	
53		63					0	94	1	1	1		3	
54		64					0	95	3	1	1	1	6	
55		65					0	96	1	2	2		5	
56		66					0	97	2	1			3	
57		67					0	98		2	1		3	
58		68					0	99	1	1			2	
59		69					0	100	1	5			6	
60		70					0	101		3	2		5	
61		71					0	102	1	2	1		4	
62		72					0	103		1			1	
63		73					0	104		3			3	
64		74	1	1			2	105		1			1	
65		75					0	106		2			2	
66		76					0	107	1			1	2	
67		77	1				1	108		2			2	
68		78		1			1	109	1				1	
69		79					0	110	1				1	
70		80		1			1	111		1			1	
71		81					0	112					0	
72		82					0	113	1				1	
73		83		3		1	4	114		2			2	
74		84					0	115		2			2	
75		85		2	1		3	116		1			1	
76		86		3			3	117					0	
77		87		1			1	118					0	
78		88			1		1	119					0	
79	1	89		1	2		3	120			2		2	
83	1	90	1	2	1	1	5	127		1			1	
								128		1			1	
2		3	15	5	2	0	25	16	35	11	3	0	65	
TOTAL ADULTS/OVIGERITY - - -									19	50	16	5	0	90
% ADULTS/OVIGERITY - - - -									21.1	55.5	17.	5.5	0	
MEAN 81		WEIGHTED MEAN ADULT LENGTH -- 97.7												
VAR. 8		VARIANCE OF THE MEAN LENGTH -- 121.1												
Full=90-100%, Hi=60-89%, Med=30-59%, Lo=1-29%														

Table 21. Carapace length measurement and percent ovigerity summary of sampled female red king crab captured in statistical area 636401 during the commercial king crab harvest, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Stat.Area 636401		Vessel All						Date 8 1-2 92					
JUVENILE		ADULT SIZE PER % OVIGERITY						ADULT SIZE PER % OVIGERITY					
mm	no.	mm Full	Hi	Med	Low	0	Sum	mm Full	Hi	Med	Low	0	Sum
50		60					0	91					0
51		61					0	92					0
52		62					0	93					0
53		63					0	94					0
54		64					0	95					0
55		65					0	96					0
56		66					0	97					0
57	1	67		1			1	98					0
58	1	68					0	99					0
59		69					0	100					0
60	1	70					0	101					0
61		71		1			1	102					0
62	1	72					0	103					0
63		73		1			1	104					0
64	3	74	1				1	105					0
65	2	75	2	1			3	106					0
66	4	76					0	107					0
67	3	77	1	1			2	108					0
68	2	78					0	109					0
69	1	79					0	110					0
70	3	80		1			1	111					0
71	2	81	1				1	112					0
72		82					0	113					0
73		83					0	114					0
74		84					0	115					0
75		85					0	116					0
76		86					0	117					0
77	1	87					0	118					0
78		88					0	119					0
79		89					0	120					0
80		90					0	121					0
								122					0
25		1	4	6	0	0	11	0	0	0	0	0	0
TOTAL ADULTS/OVIGERITY		--	--	--	--	--	--	1	4	6	0	0	11
% ADULTS/OVIGERITY		--	--	--	--	--	--	9.09	36.3	54.	0	0	
MEAN 66.3		WEIGHTED MEAN ADULT LENGTH -- 75.0											
VAR. 18.6		VARIANCE OF THE MEAN LENGTH -- 15.4											
Full=90-100%, Hi=60-89%, Med=30-59%, Lo=1-29%													




Figure 1. King crab fishing districts and sections of Statistical Area Q




Figure 2. Statistical areas for the Norton Sound red king crab fishery.

Legal male red king crab




Figure 3. Length frequency distribution and frequency of new and old carapace age condition of legal male king crab, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Male red king crab




Figure 4. Length frequency distribution of prerecruit, recruit, and postrecruit male red king crab, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Sublegal (prerecruit) red king crab




Figure 5 Length frequency distribution and frequency of new and old carapace age condition of prerecruit male king crab, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

Female red king crab




Figure 6. Length frequency distribution of female red king crab, juveniles, and adults, Norton Sound Section, Eastern Bering Sea, August 1-3, 1992.

